

BELARUS NATIONAL ACADEMY OF SCIENCES
B.I. STEPANOV's INSTITUTE OF PHYSICS

Red'kov V.M.¹

ON CONCEPT OF PARITY FOR A FERMION

Abstract

The known problem of fermion parity is considered on the base of investigating possible liner single-valued representations of spinor coverings of the extended Lorentz group. It is shown that in the frame of this theory does not exist, as separate concepts, P-parity and T-parity for a fermion, instead only some unified concept of (PT)-parity can be determined in a group-theoretical language.

PASC: 1130, 0230, 0365, 2110H

Key Words: fermion, parity, spinor group, exact linear representation,

8 pages, 17 references

¹E-mail: redkov@dragon.bas-net.by

1. On spinor space-time structure

It is well-known that the space-time vector $x^a = (t, x, y, z)$ can be identified with the explicit realization of the one-valued simplest representation of the (restricted) Lorentz group $L_+^\uparrow$. But in nature we face particles of integer and half-integer spin, bosons and fermions. Among respective sets of Lorentz group representations, entities describing bosons and fermions, there exists a clear-cut distinction: boson-based representations $T_{bos.}$ are single-valued whereas fermion-based representations $T_{ferm.}$ are double-valued on the group $L_+^\uparrow$. In other words, $T_{bos.}$ are global representations, whereas $T_{ferm.}$ are just local ones. One might have advanced a number of theoretical arguments to neglect such a light trouble. However, the fact of prime importance is that this global-local difficulty cannot be cleared up – in the frames of the orthogonal group $L_+^\uparrow = SO_0(3,1)$ it is insurmountable [1-33].

It has long been known that to work against the global-local problem one must investigate and employ one-valued representations (boson-based as well as fermion-based) of the covering group $SL(2, C)$. The group $SL(2, C)$ is other group, different from $L_+^\uparrow = SO_0(3,1)$, but it is linked to the latter by a quite definite homomorphic mapping [1-5]. At this every local representation of the orthogonal group has its counterpart – global representation of the covering group; evidently, it is a quite usual procedure.

In essence, a strong form of this changing $L_+^\uparrow = SO_0(3,1) \implies SL(2, C)$, when instead of the orthogonal Lorentz group $L_+^\uparrow = SO_0(3,1)$ we are going to employ the covering group $SL(2, C)$ and its representations throughout, and in addition we are going to work in the same manner at describing the space-time structure itself, is the known Penrose-Rindler spinor approach. Else one point should be emphasized in this context: there exists a close connection between the above spinor approach in physics and the known Kustaanheimo-Stiefel technics². Even more, one may say that these two methods are intrinsically the same, though looking differently.

An essential feature of the present work is the use of 4-spinors (bispinors in physical terms) instead of 2-spinors that are the base for the most literature. The foundation to do so is that among 4-spinors there exists real-valued ones, so-called Majorana 4-spinors [6]. The simplest real-valued 4-spinor depends upon four real parameters, exactly as in the case of Weyl complex 2-spinor. But these two models are not equivalent because all properties under spinor discrete symmetry transformations are immersed in the 4-spinor technics from the very beginning, whereas in the 2-spinor model it is not so.

Therefore, starting from real 4-spinors, as fundamental physical objects, one may expect to embed discrete symmetry operations in spinor form and simultaneously to trace in detail the appearance of the imaginary i in the model, which is of prime importance in quantum theory (for example, by translating the real Majorana wave equation into a complex Dirac's).

The present work is restricted to investigating only the problem of accurate description of the one-valued representations of four different spinor groups, all covering the full Lorentz group $L_{+-}^{\uparrow\downarrow}$, including P and T - reflections. The results obtained will provide us with the base for a new discussion of the old fermion parity problem [9-34].

2. Spinor covering of the total Lorentz group $L_{+-}^{\uparrow\downarrow}$

By adding to the set of proper orthochronous Lorentz matrices

$$L_a^b(k, k^*) = \bar{\delta}_c^b [-\delta_c^b k^n k_n^* + k_c k^{b*} + k_c^* k^b + i \epsilon_c^{bmn} k_m k_n^*], \quad L(k, k^*) = L(-k, -k^*) \quad (2.1a)$$

two matrices

$$P: L_a^{(P)b} = +\bar{\delta}_a^b, \quad T: L_a^{(T)b} = -\bar{\delta}_a^b \quad (2.1b)$$

one readily produces the total Lorentz group $L_{+-}^{\uparrow\downarrow}$. The commutation rules between $L_a^b(k, k^*)$ and the discrete elements P, T are defined by the relation

$$\bar{\delta}_a^b L_b^c(k, k^*) = L_a^b(\bar{k}^*, \bar{k}) \bar{\delta}_b^c. \quad (2.1c)$$

It is known that the group $L_{+-}^{\uparrow\downarrow}$ has four types of vector representations:

$$T_a^b(L) = f(L) L_a^b, \quad L \in L_{+-}^{\uparrow\downarrow}. \quad (2.2a)$$

²The author is grateful to Yu.A. Kurochkin for pointing out this connection and the desirability to have investigated it in more detail; also the author is grateful to E.A. Tolkachev for explanation of the main ideas of the Kustaanheimo-Stiefel approach.

Here $f(L)$ stands for a real-valued function on the group $L_{+-}^{\uparrow\downarrow}$ obeying the equation

$$f(L_1) f(L_2) = f(L_1 L_2) . \quad (2.2b)$$

Eq. (2.2b) can be satisfied by four different functions ($L \in L_{+-}^{\uparrow\downarrow}$):

$$\begin{aligned} f_1(L) &= 1 , \quad f_2(L) = \det(L) , \\ f_3(L) &= \text{sgn} (L_0^0) , \quad f_4(L) = \det(L) \text{sgn} (L_0^0) . \end{aligned} \quad (2.2c)$$

Correspondingly there exist four 4-vector representations of the full Lorentz group :

$$\begin{aligned} 1. \quad & \begin{aligned} T_1(L) &= L \\ T_1(P) &= +P \\ T_1(T) &= +T \end{aligned} & 2. \quad & \begin{aligned} T_2(L) &= L \\ T_2(P) &= -P \\ T_2(T) &= -T \end{aligned} \\ 3. \quad & \begin{aligned} T_3(L) &= L \\ T_3(P) &= +P \\ T_3(T) &= -T \end{aligned} & 4. \quad & \begin{aligned} T_4(L) &= L \\ T_4(P) &= -P \\ T_4(T) &= +T \end{aligned} \end{aligned} . \quad (2.3)$$

It is readily verified that these four are non-equivalent ones.

It should be emphasized that the above-described enlargement of the group $L_a^b(k, k^*)$ by adding two discrete operations P and T is not an enlargement of the spinor group $SL(2.C)$, instead it is only an expansion of the orthogonal group $L_-^{\uparrow}$. From a spinor view-point the operations P and T are transformations acting in the space of 2-rank spinors but not in the space of 1-rank spinors. Often, all group-theoretical consideration of discrete symmetry problems is reduced only to this treatment. Evidently, a more comprehensive study of P, T -symmetry can be done in the framework of first-rank spinors, when one extends the covering group $SL(2.C)$ by adding spinor discrete operations.

Now we can start to solve this task. A covering group for the total Lorentz group can be made constructively by adding two specific 4×4 -matrices to the known set of bispinor transformations $\tilde{SL}(2.C)$ of the group $SL(2.C)$

$$S(k, \bar{k}^*) = \begin{pmatrix} B(k) & 0 \\ 0 & B(\bar{k}^*) \end{pmatrix} \in \tilde{SL}(2.C) \sim SL(2.C). \quad (2.4)$$

Those two new matrices are to be taken from the following

$$M = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} , \quad M' = iM , \quad N = \begin{pmatrix} 0 & -iI \\ +iI & 0 \end{pmatrix} , \quad 'N = iN \quad (2.5)$$

with the multiplication table

$$\begin{array}{cccc} & M & M' & N & 'N \\ M & \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} & \begin{pmatrix} iI & 0 \\ 0 & iI \end{pmatrix} & \begin{pmatrix} +iI & 0 \\ 0 & -iI \end{pmatrix} & \begin{pmatrix} -I & 0 \\ 0 & +I \end{pmatrix} \\ M' & \begin{pmatrix} +iI & 0 \\ 0 & +iI \end{pmatrix} & \begin{pmatrix} -I & 0 \\ 0 & -I \end{pmatrix} & \begin{pmatrix} -I & 0 \\ 0 & +I \end{pmatrix} & \begin{pmatrix} -iI & 0 \\ 0 & +iI \end{pmatrix} \\ N & \begin{pmatrix} -iI & 0 \\ 0 & +iI \end{pmatrix} & \begin{pmatrix} +I & 0 \\ 0 & -I \end{pmatrix} & \begin{pmatrix} +I & 0 \\ 0 & +I \end{pmatrix} & \begin{pmatrix} +iI & 0 \\ 0 & +iI \end{pmatrix} \\ 'N & \begin{pmatrix} +I & 0 \\ 0 & -I \end{pmatrix} & \begin{pmatrix} +iI & 0 \\ 0 & -iI \end{pmatrix} & \begin{pmatrix} +iI & 0 \\ 0 & +iI \end{pmatrix} & \begin{pmatrix} -I & 0 \\ 0 & -I \end{pmatrix} \end{array} \quad (2.6)$$

It should be noted that the squared matrices $M, M', N, 'N$ equal $+I, -I \in \tilde{SL}(2.C)$. Therefore, having added any two elements from these we will have extended the group $\tilde{SL}(2.C)$ in fact by two new operations only. Also, because the group $L(2.C)$ contains $-I$, extension of the group by any two from $-M, -M', -N, -'N$ results in the same.

However, if one chooses any other different from $+1, -1, +i, -i$ phase factors at $M, M', N, 'N$, then one will obtain substantially new extended groups. For instance, addition of the discrete element αM with $\alpha = \exp(i2\pi/n)$ leads to a group extended by $(n-1)$ new elements:

$$\alpha M, \alpha^2 I, \alpha^3 M, \alpha^4, \dots, \alpha^{(n-1)} M.$$

Those will not be considered in the present work.

So, enlarging the group $\tilde{SL}(2.C)$ as was shown above, we obtain six covering groups:

$$\begin{aligned} G_M &= \{ S(k, \bar{k}^*) \uplus M \uplus M' \}, & G_N &= \{ S(k, \bar{k}^*) \uplus N \uplus 'N \}, \\ G' &= \{ S(k, \bar{k}^*) \uplus M' \uplus N \}, & 'G &= \{ S(k, \bar{k}^*) \uplus 'N \uplus M \}, \\ G &= \{ S(k, \bar{k}^*) \uplus M \uplus N \}, & 'G' &= \{ S(k, \bar{k}^*) \uplus M' \uplus 'N \} \end{aligned} \quad (2.7)$$

with multiplication tables

$$\begin{aligned} G_M : \quad M^2 &= +I, & M'^2 &= -I, & MM' &= M'M; \\ G_N : \quad N^2 &= +I, & 'N^2 &= -I, & N'N &= 'NN; \\ G' : \quad M'^2 &= -I, & N^2 &= +I, & M'N &= -NM'; \\ 'G : \quad 'N^2 &= -I, & M^2 &= +I, & 'NM &= -M'N; \\ G : \quad M^2 &= +I, & N^2 &= +I, & MN &= -NM; \\ 'G' : \quad M'^2 &= -I, & 'N'^2 &= -I, & M' 'N &= -'NM' \end{aligned} \quad (2.8)$$

and

$$F S(k, \bar{k}^*) = S(\bar{k}^*, k) F, \quad F \in \{M, M', N, 'N\}. \quad (2.9)$$

One can notice that the tables for the groups G_M and G_N happen to coincide; as well those of G' and $'G$. This implies that the groups G_M and G_N represent the same abstract group, as well as G and $'G$. Indeed, it is readily verified that G_M and G_N , as well as G' and $'G$, can be inverted into each other by a similarity transformation:

$$G_N = A G_M A^{-1} : \quad A S(k, \bar{k}^*) = S(\bar{k}^*, k) A, \quad (2.10a)$$

$$A M A^{-1} = +N, \quad A M' A^{-1} = +'N,$$

$$A = \text{const} \begin{pmatrix} -iI & 0 \\ 0 & +I \end{pmatrix};$$

$$'G = A G' A^{-1} : \quad A S(k, \bar{k}^*) = S(\bar{k}^*, k) A, \quad (2.10b)$$

$$A M' A^{-1} = +'N, \quad A N A^{-1} = -M,$$

$$A = \text{const} \begin{pmatrix} -iI & 0 \\ 0 & +I \end{pmatrix}.$$

In other words, above only four different covering groups were defined. As long as in literature all six variants are encountered, all six will be traced below.

3. Representations of extended spinor groups

Now let us construct exact linear representations of the groups $G_M, G_N, G', 'G, G, 'G'$. It suffices to consider in detail only one group, for definiteness let it be G_M . Its multiplication table is

$$M^2 = -I, \quad M'^2 = -I, \quad M M' = M M \quad (3.1)$$

$$F S(k, \bar{k}^*) = S(\bar{k}^*, k) F, \quad (F = M, M'),$$

$$(k_1, \bar{k}_1^*)(k_2, \bar{k}_2^*) = (< k_1, k_2 >, < \bar{k}_1^*, \bar{k}_2^* >).$$

Here the symbol $<, >$ stands for the known multiplication rule for the vector-parameter on the group $SL(2.C)$:

$$< k_1, k_2 > = (k_1^0 k_2^0 + \vec{k}_1 \vec{k}_2; \vec{k}_1 k_2^0 + k_2^0 \vec{k}_1 + i [\vec{k}_1 \vec{k}_2]).$$

Let us look for solution of the problem $g \longrightarrow T(g)$ in the form

$$T(g) = f(g) g, \quad g \in G_M, \quad f(g_1) f(g_2) = f(g_1 g_2) \quad (3.2a)$$

where $f(g)$ is a numerical function on the group G_M . Substituting (3.2a) into (3.1) we get to

$$\begin{aligned} [f(M)]^2 &= f(I) , \quad [f(M')]^2 = f(-I) , \\ f(S(k, \bar{k}^*)) &= f(S(\bar{k}^*, k)) , \\ f(S(k_1, \vec{k}_1^*)) f(S(k_2, \vec{k}_2^*)) &= f(S(< k_1, k_2 > , < \vec{k}_1^* , \vec{k}_2^* >)) . \end{aligned}$$

There exist four different functions f_i satisfying the equation above:

$$\begin{array}{ccccc} G_M & f_1(g) = & f_2(g) = & f_3(g) = & f_4(g) = \\ S(k, \bar{k}^*) & +1 & +1 & +1 & +1 \\ M & +1 & -1 & +1 & -1 \\ M' & +1 & -1 & -1 & +1 . \end{array} \quad (3.2b)$$

Correspondingly we have four representations $T_i(g)$ of the group G_M . In the same manner one can construct analogous representation $T_i(g)$ of remaining five groups. All those can be described by the following table

$$\begin{array}{cccccc} g = & T_1(g) = & T_2(g) = & T_3(g) = & T_4(g) = & \\ & S(k, \bar{k}^*) & S(k, \bar{k}^*) & S(k, \bar{k}^*) & S(k, \bar{k}^*) & S(k, \bar{k}^*) \\ G_M & M & +M & -M & +M & -M \\ & M' & +M' & -M' & -M' & +M' \\ G_N & N & +N & -N & +N & -N \\ & 'N & +'N & -'N & -'N & +'N \\ G' & M' & +M' & -M' & +M' & -M' \\ & N & +N & -N & -N & +N \\ 'G & 'N & +'N & -'N & +'N & -'N \\ & M & +M & -M & -M & +M \\ G & M & +M & -M & +M & -M \\ & N & +N & -N & +N & -N \\ 'G' & M' & +M' & -M' & +M' & -M' \\ & 'N & +'N & -'N & -'N & +'N \end{array} \quad (3.3)$$

For each of the groups under consideration one can ask a question: are the four representations $T_i(g)$ equivalent or not. With the help of relations

$$F = \text{const} \begin{pmatrix} -I & 0 \\ 0 & +I \end{pmatrix} , \quad F S(k, \bar{k}^*) F^{-1} = S(k, \bar{k}^*) , \quad (3.4a)$$

$$F M F^{-1} = -M , \quad F M' F^{-1} = -M' F ,$$

$$N F^{-1} = -N , \quad F 'N F^{-1} = -'N$$

it is easily follows that the type $T_2(g)$ is equivalent to the type $T_1(g)$, as well as $T_4(g)$ is equivalent to $T_3(g)$:

$$T_2(g) = F T_1(g) F^{-1} , \quad T_4(g) = F T_3(g) F^{-1} . \quad (3.4b)$$

Summarizing, we have got to the following: for every of six groups only two non-equivalent representations $g \rightarrow T(g) = f(g) g$ are possible: $T_1(g) \sim T_2(g)$ $T_3(g) \sim T_4(g)$. Evidently, this result does not depend on an explicit realization of the discrete spinor transformations. The above study of the exact linear representations of the extended spinor groups leads to a new concept of a space-time intrinsic parity of a fermion. In group-theoretical terms P -parity and T -parity do not have any sense, instead only their joint characteristic, that might be called (PT) -parity, can be defined in the group-theory framework³.

³It should be added that a very similar fact was noted by Yu. Shirokov [28] and De Vitt [32].

4. Representations of the coverings for $L_{+-}^{\uparrow}$ and $L_{+-}^{\uparrow\downarrow}$

Now we are going to consider the problem of linear representations of the spinor groups that are supposedly cover the partly extended Lorentz groups $L_{+-}^{\uparrow}$ and $L_{+-}^{\uparrow\downarrow}$ (improper orthochronous and proper non-orthochronous respectively). Such coverings of partly extended groups can be constructed by adding any one matrix from M, M', N, N' .

It is known that orthogonal groups $L_{+-}^{\uparrow}$ and $L_{+-}^{\uparrow\downarrow}$ have vector representations of only two types. The case of the group $L_{+-}^{\uparrow}$:

$$T_1 = T_3 : L \implies L = (\text{sgn } L_0^0) L , \quad (4.1a)$$

$$T_2 = T_4 : L \implies L = (\det L) L = (\det L)(\text{sgn } L_0^0) L . \quad (4.1b)$$

The case of the group $L_{+-}^{\uparrow\downarrow}$ looks as

$$T_1 = T_4 : L \implies L = (\det L)(\text{sgn } L_0^0) L , \quad (4.2a)$$

$$T_2 = T_3 : \implies L = (\det L) L = (\text{sgn } L_0^0) L . \quad (4.2b)$$

Now let us proceed to partly extended spinor groups, coverings for $L_{+-}^{\uparrow\downarrow}$ and $L_{+-}^{\uparrow}$. With the use of one additional discrete operation one can determine four extended spinor groups:

$$\tilde{SL}(2.C)_M = \{ S(k, \bar{k}^*) \oplus M \} \text{ and so on .} \quad (4.3)$$

In so doing extended groups $\tilde{SL}(2.C)_M$, $\tilde{SL}(2.C)_N$ turn out to be isomorphic. Analogously, $\tilde{SL}(2.C)_{M'}$ is isomorphic to $\tilde{SL}(2.C)_{N'}$. Thus, there exist only two different spinor groups, each of them covers both $L_{+-}^{\uparrow\downarrow}$ and $L_{+-}^{\uparrow}$:

$$\tilde{SL}(2.C)_M \sim \tilde{SL}(2.C)_N , \quad \tilde{SL}(2.C)_{M'} \sim \tilde{SL}(2.C)_{N'} .$$

Now, we are to list simplest representations of these groups. The result obtained is as follows: all above representations $T_i(g)$ (see Section 3) at confining them to sub-groups $SL(2.C)_{M(N)}$ and $SL(2.C)_{M',(N')}$ lead to representations changing into each other by a similarity transformation. In other words, in fact there exists only one representation of these partly extended spinor groups. This may be understood as impossibility to determine any group-theoretical parity concept (P or T) within the limits of partly extended spinor groups.

5. On reducing spinor groups to a real form

Till now we have counted all spinor groups $G_M \sim G_N$, $G' \sim' G$, G , G' among possible candidates to be covering of the full Lorentz group $L_{+-}^{\uparrow\downarrow}$. It is desirable to have formulated some extra argument to choose only one spinor group as a genuine (physical) covering.

Let us draw attention to the fact that the used bispinor matrix (continuous one)

$$\Phi = \begin{pmatrix} \xi^\alpha \\ \eta_{\dot{\alpha}} \end{pmatrix} ; \quad \Phi' = S(k, \bar{k}') \Phi , \quad S = \begin{pmatrix} B(k) & 0 \\ 0 & B(\bar{k}^*) \end{pmatrix} \quad (5.1)$$

can be taken to a real-valued form. This implies that in the bispinor space a special basis can be found where the bispinor wave function

$$\Phi_M(x) = \varphi(x) + i\xi(x) \quad \varphi^*(x) = \varphi(x) , \quad \xi^*(x) = \xi(x)$$

transforms under $SL(2.C)$ group by means of real (4×4) -matrices. Therefore, the real 4-spinors $\varphi(x)$ and $\xi(x)$, constituents of complex-valued $\Phi_M(x)$, transform as independent irreducible representations. In physical context of real Majorana fermions this reads as a group-theoretical permission to exist. But these arguments have been based only on continuous $SL(2.C)$ -transformations, the idea is to extend these on discrete operations too.

So we must find the answer to the question of which of the extended spinor groups can be reduced to a real form? With this end in mind let us write down the bispinor matrix in the form that does not depend on an accidental basis choice⁴:

$$S(k, \bar{k}^*) = \frac{1}{2}(k_0 + k_0^*) + \frac{1}{2}(k_0 - k_0^*) \gamma^5 + (k_1 + k_1^*) \sigma^{01} + (k_1 - k_1^*) i \sigma^{23} + \\ (k_2 + k_2^*) \sigma^{02} + (k_2 - k_2^*) i \sigma^{31} + (k_3 + k_3^*) \sigma^{03} + (k_3 - k_3^*) i \sigma^{13} . \quad (5.2)$$

⁴ Above we employed the Weyl basis.

The form (5.2) being taken in the spinor frame

$$\gamma^a = \begin{pmatrix} 0 & \bar{\sigma}^a \\ \sigma^a & 0 \end{pmatrix}, \quad \gamma^5 = -i \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} -I & 0 \\ 0 & +I \end{pmatrix} \quad (5.3)$$

gives the used above. But the form (5.2) is invariant under any similarity transformations:

$$\gamma^{a'} = A \gamma^a A^{-1}, \quad \Phi' = A \Phi, \quad S'(k, \bar{k}^*) = A S(k, \bar{k}^*) A^{-1}.$$

So, it remains to write down all used discrete operations $M, M'N, 'N$ in terms of Dirac matrices:

$$M = +\gamma^0, \quad M' = +i \gamma^0, \quad N = +i \gamma^5 \gamma^0, \quad 'N = -\gamma^5 \gamma^0 \quad (5.4)$$

and to take into account that any Majorana basis satisfy the relations

$$(\gamma_M^a)^* = -\gamma_M^a, \quad (\gamma_M^5)^* = -\gamma_M^5, \quad (\sigma_M^{ab})^* = \sigma_M^{ab}. \quad (5.5)$$

Therefore, in such frames the used discrete operations obey

$$S^* = S, \quad M^* = -M, \quad (M')^* = +M', \quad N^* = -N, \quad ('N)^* = +'N. \quad (5.6)$$

Thus, six spinor groups behave under complex conjugation as indicated below

G_M	G_N	G'	$'G$	G	$'G'$
$S^* = S$	$S^* = S$	$S^* = S$	$S^* = S$	$S^* = S$	$S^* = S$
$M^* = -M$	$N^* = -N$	$M'^* = +M'$	$'N^* = +'N$	$M^* = -M$	$M'^* = +M$
$(M')^* = +M'$	$'N^* = +'N$	$N^* = -N$	$M^* = -M$	$N^* = -N$	$'M^* = +'M$

From this it follows that only the group $'G'$ can be reduced to a real-valued form. Only this group allows for real-valued spinor representations, Majorana fermions⁵.

6. Discussion

In the present paper, the problem of fermion parity is considered on the base of investigating possible single-valued representations of spinor coverings of the extended Lorentz group. It is shown that in the frame of this theory does not exist, as separate concepts, P-parity and T-parity for a fermion, instead only some unified concept of (PT)-parity can be determined in a group-theoretical Apparently, physics with spinor group in its base differs from that based on the orthogonal group $L_{+-}^{\uparrow\downarrow}$ and only experiment must decide the problem once and for all. It is needless to say that the task cannot be solved without a thorough theoretical analysis of possible experimental verifications in both orthogonal and spinor framework. language.

References

1. Lipschitz R. Bonn, 1886.
2. Darboux G. // Bull. des Sciences Math. 1905, Tom 9, 2 série (Ce memoire est un résumé des leçons que l'auteur a faites à la Sorbonne en 1900 et 1904.).
3. Cartan E. // Bull. Soc. Math. France. 1913, Tom 41, P. 53-96.
4. Weyl H. // Math. Z. 1924, Bd 23, S. 271-309; 1925, Bd 24, S. 329-395.
5. Schreier O. // Hamb. Abh. Math. Sem. 1925, Bd. 4, S. 15-32; // Hamb. Abh. Math. Sem. 1926, Bd. 5, S. 236-244.
6. Majorana E. // Nuovo Cim. 1937, Vol. 14, P. 171-184.
7. Racah G. Sulla simmetria tra particelle e antiparticell. // Nuovo Cim. 1937, Vol. 14, P. 322.
8. Cartan E. // Actualités Sci. et Ind. 1938, n. 643, Hermann, Paris; 1938, n. 701, Hermann, Paris.
9. Yang C.N., Tiomno J. // Phys. Rev. 1950. Vol. 79, n. 3, P. 495-498.
10. Jarkov G.F. JETP, 1950, Vol. 20, No 6, P. 492-496.
11. Caianiello E.R. // Nuovo Cim. 1951, Vol. 7, P. 534; 1951, Vol. 8, P. 749; 1952, Vol. 9, P. 336.
12. Wick G.C., Wigner E.P., Wightman A.S. // Phys. Rev. 1952, Vol. 88, P. 101.

⁵This variant coincides with the known in the literature Racah group [7].

13. Shapiro I.C. // JETP. 1952, Vol. 22, No 5, P. 524-538.
14. Shapiro I.C. // UFN. 1954, Vol. 53, No 1, P. 7-68.
15. Watanabe S. // Rev. Mod. Phys. 1955, Vol. 27, P. 26-39; 1955, Vol. 27, P. 40-76.
16. Heine V. // Phys. Rev. 1957, Vol. 107, n. 2, P. 620-623.
17. Широков Ю.М. // ЖЭТФ. 1957, Том 33, Вып. 4, С. 861-872; 1957, Том 33, Вып. 5, С. 1196-1207; 1957, Том 33, Вып. 5, С. 1208-1214; 1958, Том 34, Вып. 3, С. 717-724; 1959, Том 36, Вып. 3, С. 879-888.
18. Feinberg G., Weinberg S. // Nuovo Cim. 1959, Vol. 14, n. 3, P. 571-592.
19. Широков Ю.М. // Nucl. Phys. 1960, Vol. 15, P. 1-12; 1960, Vol. 15, P. 13-15.
20. Богущ А.А., Федоров Ф.И. // Докл. АН БССР. 1961, Том 5, n. 8, С. 327-330.
21. Sudarshan E.C.G. // J. Math. Phys. 1965, Vol. 6, n. 8, P. 1329-1331.
22. Macfarlane A.J. // J. Math. Phys. 1962, Vol. 3, P. 1116-1129.
23. Wigner E. In: Group theoretical concepts and methods in elementary particle physics. Ed. Gürsey F.N.Y., 1964, Lectures at the Istanbul Summer School of theoretical physics. 1962, P. 37-80.
24. Lee T.D. // Phys. Today. 1966, Vol. 19, n. 3, P. 23-31.
25. Lee T.D., Wick G.C. // Phys. Rev. 1966, Vol. 148, n. 4, P. 1385-1404.
26. Goldberg H. // Nuovo Cim. A. 1969, Vol. 60, n. 4, P. 509-518.
27. Wigner E. Symmetry and reflections. Indiana University Press, Bloomington - London, 1970.
28. Широков Ю.М. // ЭЧАЯ. 1972 Том 3, Вып 3, С. 606-649; 1973. Том 4, Вып. 1, С. 42-78.
29. Yang C.N. // J. de Physique. 1982, Colloque C8, P. 439.
30. Altmann S.L. Rotations, quaternions, and double groups. 1986.
31. В.С. Де Витт. Динамическая теория групп и полей. Москва, Наука, 1987.
32. Sharma C.S. // Nuovo Cim. B. 1989, Vol. 103, P. 431-434.
33. Recai Erdem. 12 pages; **hep-th/9809054 V2**